



Research Paper

Obesity Gene Atlas in Mammals

Tanja Kunej^{1*}, Dasa Jevsinek Skok^{1*}, Minja Zorc^{1,2*}, Ana Ogrinc¹, Jennifer J. Michal³, Milena Kovac¹, and Zhihua Jiang^{3*}

1. Department of Animal Science, Biotechnical Faculty, University of Ljubljana, Domzale, Slovenia;
2. Centre for Mathematical and Computational Biology, Rothamsted Research, Harpenden, Herts, AL5, 2JQ, UK;
3. Department of Animal Sciences, Washington State University, Pullman, Washington 99164-6351, USA.

* Authors contributed equally to the work.

✉ Corresponding author: Dr. T. Kunej, Department of Animal Science, Biotechnical Faculty, University of Ljubljana, Domzale, Slovenia, Tel: +386 13203890; FAX: +386 17241005; E-mail address: tanja.kunej@bf.uni-lj.si and Prof. Zhihua Jiang, Department of Animal Sciences, Washington State University, Pullman, Washington 99164-6351, E-mail address: jiangz@wsu.edu.

© Ivyspring International Publisher. This is an open-access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by-nc-nd/3.0/>). Reproduction is permitted for personal, noncommercial use, provided that the article is in whole, unmodified, and properly cited.

Published: 2012.02.10

Abstract

Obesity in humans has increased at an alarming rate over the past two decades and has become one of the leading public health problems worldwide. Studies have revealed a large number of genes/markers that are associated with obesity and/or obesity-related phenotypes, indicating an urgent need to develop a central database for helping the community understand the genetic complexity of obesity. In the present study, we collected a total of 1,736 obesity associated loci and created a freely available obesity database, including 1,515 protein-coding genes and 221 microRNAs (miRNAs) collected from four mammalian species: human, cattle, rat, and mouse. These loci were integrated as orthologs on comparative genomic views in human, cattle, and mouse. The database and genomic views are freely available online at: http://www.integratomics-time.com/fat_deposition. Bioinformatics analyses of the collected data revealed some potential novel obesity related molecular markers which represent focal points for testing more targeted hypotheses and designing experiments for further studies. We believe that this centralized database on obesity and adipogenesis will facilitate development of comparative systems biology approaches to address this important health issue in human and their potential applications in animals.

Key words: adipogenesis, fat deposition, integratomics, mammals, microRNA (miRNA), obesity.

Introduction

Obesity is a result of excess body fat accumulation, which often causes adverse effects on human health, such as cardiovascular diseases, type 2 diabetes mellitus, and cancer. Depending on suspected etiology, obesity is commonly classified into three subgroups: monogenic, syndromic, and polygenic or common obesity [1]. Incidence of obesity has dramatically increased over the past few decades and is a global health problem worldwide. In particular, polygenic obesity depends on an individual's genetic

makeup that is susceptible to an environment that promotes energy consumption over energy expenditure. As such, search for genes associated with polygenic obesity has focused on multiple interacting alleles contributing to common diseases. Major approaches used to identify novel gene variants associated with polygenic obesity, include candidate gene, genome-wide linkage and genome-wide association studies [2]. Genes that define polygenic obesity have been implicated in a wide variety of biological

functions, such as the regulation of food intake, energy expenditure, lipid and glucose metabolism, and adipose tissue development [3]. There is growing evidence for a relationship between obesity-associated insulin resistance and mitochondrial dysfunction [4].

Characterization of genes responsible for increased fat deposition is needed to develop methods and tools that can identify people at high risk for obesity. This information can also be used to design appropriate targeted therapies that will improve the quality of care of obesity-linked conditions [5]. The identification of obesity candidate loci is also attractive for animal breeding, due to growing consumer demand for products with lower fat content. On the other hand, selection of animals with superior marbling or intramuscular fat (IMF) phenotypes is an important objective for livestock producers since meat with a high marbling content tends to be juicier and is more tender and flavorful than very lean meat [6].

The rapid development of molecular genetic marker technology in recent years has led to the identification of genes that contribute to the genetic variation (so called quantitative trait loci, QTL) of marbling or IMF content in livestock species, and hence to molecular farming by marker-assisted selection [7, 8]. Therefore, the same orthologous genes may have conserved functions in biological or biochemical pathways, and thus explain the same or similar variations of the concordant QTL among different species [9]. Comparative approach may reveal novel candidate genes and functional insights into obesity in human [10]. In a previous study, we collected over 2,000 reports on genes/markers affecting fat phenotypes in several species [5], assigned them to the human orthologous regions and subsequently used the markers for identification of genetic networks associated with various fat and fat-related phenotypes in a Wagyu x Limousin cattle population [11].

Recent discoveries also link the development of obesity to microRNAs (miRNAs) [12]. MiRNAs regulate expression of most genes and play critical roles in different biological processes, such as cell differentiation, proliferation, death, metabolism, and energy homeostasis [13]. Some miRNAs have also been implicated in regulation of adipogenic differentiation [14-27]. These data strongly suggest that miRNAs represent a new class of adipogenic inhibitors that may play a role in the pathological development of obesity [19].

The human obesity gene map was updated yearly from 1996 to 2006 and published in the journal *Obesity* (previously, *Obesity Research*). This map displayed gene/marker information associated with

obesity and obesity-related phenotypes using data collected from a variety of sources, such as PubMed using a combination of key words, authors, and journals; continuous reviews of obesity and genetics journals; personal collection of reprints; and papers made available to the authors by colleagues around the world [8]. Although the yearly review was a valuable tool for researchers involved in this field, it was discontinued after 2006. However, an increasing number of studies have been published on obesity and adipogenesis in human as well as in other animals.

The aim of the present study was, therefore, to assemble all available information associated with fat deposition into a publicly available online obesity database. We collected data with obesity associated genomic loci from different sources and species (human, cattle, rat, and mice) in an obesity gene atlas, which includes a database and genomic views. We believe that this database will serve as a valuable source for fine mapping and narrowing QTL regions to a few genes and identification of major pathways involved in obesity. In order to integrate data from different sources, a holistic (map-driven) approach was used and an interactive genomic view of collected obesity candidate loci was developed. The map-based approach reveals positional overlaps between candidate loci, thus allowing complementation of different pieces of information from various species for cloning of positional candidate genes.

Material and Methods

Obesity associated candidate loci identified in human, cattle, rat, and mice were collected from literature and publicly available databases. All relevant publications were identified after searching PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) and Web of Science (<http://apps.webofknowledge.com/>), with key phrases, such as gene, genetics, epigenetics, non-coding RNA, microRNA, obesity, adipose tissue, marbling, fat deposition, adipogenesis, human, cattle, rat, and mouse. Obesity-associated loci were also extracted from the following databases: Human obesity gene map (OGM) [8], GeneCards, version 3.07 (<http://www.genecards.org>), Mouse Genome Informatics, release 4.42 (MGI: <http://www.informatics.jax.org>). To analyze genomic overlaps of obesity associated loci, QTL were downloaded from Rat Genome Database, release date: October 2011 (RGD: <http://rgd.mcw.edu/>) [28], and Animal QTL Database, release 15 (<http://www.animalgenome.org/cgi-bin/QTLdb/index/>). QTL associated with obese phenotypes were selected (body fat, body weight, adiposity, obesity, and diabetes) for cattle,

mouse, and human. Gene nomenclature was based on the HUGO Gene Nomenclature Guidelines (<http://www.genenames.org>). MiRNA nomenclature was adjusted according to the miRBase, release 18 (<http://microrna.sanger.ac.uk/>). Ingenuity Pathway Analysis (IPA: Ingenuity Systems® www.ingenuity.com) was used to interpret the obesity candidate genes in the context of biological processes and pathways. The Ingenuity Knowledge Base contains information from databases including: Gene Ontology, Entrez Gene, RefSeq, OMIM, KEGG metabolic pathway information etc. (www.ingenuity.com). IPA analysis was performed twice, once with protein coding genes and miRNAs and the second time only with protein coding genes. Genomic views (graphical overview of the chromosomal locations) of obesity associated loci were constructed using the web-based

interactive visualization tool Flash GViewer (<http://gmod.org/wiki/Flashgviewer/>) developed by the GMOD project.

Results and Discussion

Collecting obesity related data and creating the database

Previous studies have confirmed a large number of genetic loci associated with obesity and adipogenesis in mammals. We retrieved data from the literature and publicly available databases (PubMed, Web of Science, OGM, GeneCards, MGI, and RGD) for 1,515 protein-coding genes and 221 miRNAs in four different species (human, cattle, rat, and mouse). The workflow of data collection is presented in **Figure 1**.




Figure 1: The workflow of data collection. A total of 1,736 loci were retrieved from five databases and current literature for four mammalian species (Rat, Human, Cattle, and Mouse). Genes and miRNAs associated with adipogenesis were retrieved from literature, OGM, Gene Cards, MGI, and RGD. OGM: Obesity gene map, RGD: Rat genome database, MGI: Mouse genome informatics database.

As early as in 2004, miRNAs were identified to play regulatory roles in adipocyte differentiation [14]. Since then, 221 miRNAs have been found to be expressed or dysregulated in adipose tissue or adipocytes in human, cattle, rat, and mouse. Those miRNAs were collected in studies using 3T3-L1 pre-adipocytes, adipocytes, adipogenesis, adipose tissue, back fat, diabetes, omental fat and subcutaneous fat, and obesity (Figure 2) [14-27]. As shown on a Venn diagram (Figure 3), some miRNAs have been found to be as-

sociated with fat deposition in more than one species. For example, 14 miRNAs (let-7a, let-7b, let-7c, let-7e, let-7f, mir-103, mir-10b, mir-125a, mir-125b, mir-143, mir-23a, mir-23b, mir-26a, and mir-99b) have been reported to affect fat deposition in all four species. Interestingly, Wang et al. [29] also described involvement of mir-143 in regulation of porcine adipocyte lipid metabolism, which indicates its strong impact on adipogenesis.




Figure 2: (A) MiRNAs associated with 3T3-L1 pre-adipocytes, adipocytes, adipogenesis, adipose tissue, back fat, diabetes, omental fat and subcutaneous fat, and obesity retrieved from current literature. An example for miRNA let-7b associated with five obesity-associated traits is shown in red. (B) Close up of miRNAs with regulatory roles in obesity-associated traits. (C) MiRNAs associated with back fat.




Figure 3: Venn diagram presenting common obesity miRNAs for four species: cattle, human, mouse, and rat.

Based on the data collected from four mammalian species: human, mouse, rat, and cattle, we created an online database to share the information with the community on genetics and genomics of fat deposition/obesity. The database consists of 1,736 obesity-associated loci, including 1,515 protein-coding genes and 221 miRNAs. The database is freely available on the Web site at http://www.integratomics-time.com/fat_deposition. In order to further strengthen the community's involvement in the database expansion and utilization, we also generated an online data entry interface, which will enable users to update or submit new obesity-associated candidate genes and markers. The newly submitted information will be reviewed by curators before releasing to the public (http://www.integratomics-time.com/fat_deposition/add_genes).

The centralized database on genes/markers associated with obesity and adipogenesis will provide fundamental information on what genetic loci are involved in obesity-related traits, how their effects are similar or different among different species and how they interact under different conditions. The knowledge generated from the genetics/genomics studies will provide insight into the molecular basis of obesity and allow for prospective identification of people who have a genetic predisposition to become overweight and/or obese. We believe that the newly constructed database will advance us to the forefront of mapping of QTL for complex traits and understanding their biological pathways by combining a

traditional quantitative genetics approach with a modern molecular genetics approach. More importantly, the information obtained in the present study will offer potential and exciting possibilities for future development of successful therapies and new treatments for obesity in humans. On the other hand, due to growing consumer demand for products with lower fat content, an important objective in animal breeding is to improve meat quality traits. Therefore, addition of data related to livestock animal models in our database will have implications in agriculture and biomedicine.

Comparative genomic distribution of fat deposition associated loci

An integrative, comparative-genomics approach allowed us to join obesity-associated information for various species regardless of the study approach, and to present collected loci from human and animals in the form of a single species (human, mouse, and cattle) genomic view available at http://www.integratomics-time.com/fat_deposition/genomic_view/. Genomic view of obesity-associated loci is visible through Flash GViewer provided by the GMOD consortium. Human obesity associated orthologous protein-coding genes, miRNA genes and overlapping QTL are shown in **Figure 4**.

Additionally, a large number of obesity QTL have been identified in domestic and model animal species. To analyze genomic overlaps of obesity associated loci, QTL for cattle, mouse, and human were

downloaded from AnimalQTLdb and RGD. QTL associated with obese phenotype were selected: 469 in human, 355 in cattle and 274 in mouse. Genomic overlap analysis was performed comparing the location of obesity QTL, protein-coding and miRNA genes. Obesity associated orthologs overlapped with 222 human, 176 cattle, and 38 mouse QTL, respectively.

Some candidate genes overlapped with a high number of QTL and therefore could potentially be stronger molecular markers. Three human obesity candidate genes: *CEBPB* (CCAAT/enhancer binding protein (C/EBP), beta), *PTPN1* (protein tyrosine phosphatase, non-receptor type 1), and *SLPI* (secretory leukocyte peptidase inhibitor) each overlapped with 11 different QTL. Gene *ALB* (albumin) overlapped with 16 QTL in cattle. Similarly, using integratomics approach, Kunej *et al.* [25] identified a molecular marker residing within the *mmu-mir-717* gene, growth rate associated gene *Gpc3*, and growth associated QTL.

Several studies have also shown that protein-coding host genes are functionally linked with their resident miRNAs [23, 30]. MiRNA genes and their sense oriented host genes can be transcribed from shared promoters [31], whereas antisense orientation suggested that miRNAs and host genes have independent transcription mechanisms [32]. Our analysis showed that among 221 obesity-associated

miRNAs, 54 resided within human and 57 within mouse host genes, including some of non-protein-coding genes; large intergenic non-coding RNAs (lincRNAs) (**Supplementary Material: Table S1**). Sense oriented obesity associated miRNA genes from our obesity database overlapped with introns (39 in human and mouse), exons (eight in human and nine in mouse), or 3'-UTRs (two in mouse) of their host genes. Interestingly, three of the host genes have been previously associated with obesity: *mir-335* resided within host gene *MEST* (mesoderm specific transcript homolog (mouse)) in human and mouse, *mir-378* within *Ppargc1b* (peroxisome proliferator-activated receptor gamma, coactivator 1 beta), and *mir-33* within *Srebf2* (sterol regulatory element binding transcription factor 2). In addition to these three miRNA-host gene pairs, other miRNA-host gene pairs warrant further experimental analyses to explore their potential functional link.

Comparative genomics allowed exploitation of animal models for elucidation of obesity phenotype in human. However, extrapolating the gained knowledge from one species to another is often difficult, due to different anatomical and physiological characteristics. Similarly, integratomics/genomic overlap approach was already successfully used for identification of candidate loci for mammary gland associated phenotypes [33] and male infertility [34].




Figure 4: Human chromosomes with obesity-associated loci. Genomic view of the obesity-associated candidate loci presented as human orthologs. Enlargement of the chromosome 21 is showing an overlap between QTL for body weight (BW276_H) and seven obesity related loci *ADAMTS1*, *APP*, *GABPA*, *HSPA13*, *LIPI*, *NR1P1*, and *hsa-mir-99a*.

Network analysis of obesity associated genes

Human orthologs of obesity candidate genes were integrated into interactome networks using Ingenuity Pathway Analysis (IPA) (Figure 5). Input data set consisted of 1,676 genes (protein coding and miRNA genes) and can be accessed on the web page http://www.integratome.com/fat_deposition/genomic_view/. Five of the highest scored networks, diseases and disorders, molecular and cellular functions, as well as canonical pathways are shown in Figure 5a. The analysis revealed that obesity candidate genes were associated with nutritional disease, genetic, gastrointestinal, developmental disorder, and cancer. From the total of 25 top networks four of them were found to be associated with lipid metabolism (data not shown). A merged diagram of these four networks includes 140 obesity candidate genes and 175 molecules (Figure 5b, Supplementary Material: Table S2). Peroxisome proliferator-activated receptor

alpha (PPAR α) and retinoid X receptor alpha (RXR α) were identified as central nodes, also called hub molecules. Peroxisome proliferator-activated receptors (PPARs) are nuclear hormone receptors that regulate genes in energy metabolism. Activity of PPARs is dependent on RXR biological activity; PPARs require cognate lipid ligands, heterodimerization with retinoid X receptors (RXR) and coactivation by coactivators [35]. The PPAR-RXR transcriptional complex plays a critical role in energy balance, including triglyceride metabolism, fatty acid handling and storage, and glucose homeostasis: processes whose dysregulation characterize obesity, diabetes, and atherosclerosis [36]. Interestingly, our network analysis of obesity associated genes identifies both of them as central nodes. Moreover, the *PPAR α /RXR α* activation pathway was one of the top five canonical pathways identified using IPA analysis.

A

Top Networks		
ID	Associated Network Functions	Score
1	View Behavior, Digestive System Development and Function, Cell Signaling	44
2	View Lipid Metabolism, Small Molecule Biochemistry, Vitamin and Mineral Metabolism	39
3	View Cell-To-Cell Signaling and Interaction, Cellular Growth and Proliferation, Hematological System Development and Function	36
4	View Nucleic Acid Metabolism, Small Molecule Biochemistry, Genetic Disorder	35
5	View DNA Replication, Recombination, and Repair, Cellular Growth and Proliferation, Reproductive System Disease	35

Top Bio Functions		
Diseases and Disorders		
Name	p-value	# Molecules
Nutritional Disease	1,10E-243 - 1,92E-45	378
Genetic Disorder	2,41E-191 - 8,40E-36	1211
Developmental Disorder	7,35E-149 - 8,38E-40	540
Cancer	8,64E-146 - 8,18E-37	841
Gastrointestinal Disease	2,84E-118 - 2,15E-35	862

Molecular and Cellular Functions		
Name	p-value	# Molecules
Lipid Metabolism	1,90E-211 - 1,61E-35	632
Molecular Transport	1,90E-211 - 2,60E-35	778
Small Molecule Biochemistry	1,90E-211 - 3,36E-35	815
Cellular Growth and Proliferation	1,29E-164 - 5,10E-35	780
Cellular Movement	8,21E-156 - 8,90E-36	558

Top Canonical Pathways		
Name	p-value	Ratio
Leptin Signaling in Obesity	5,96E-70	71/84 (0,845)
G-Protein Coupled Receptor Signaling	1,82E-58	174/530 (0,328)
PPARα/RXRα Activation	1,3E-57	97/186 (0,522)
Hepatic Cholestasis	3,63E-52	84/176 (0,477)
Role of Macrophages, Fibroblasts and Endothelial Cells in Rheumatoid Arthritis	2,63E-50	122/333 (0,366)




Figure 6: Canonical pathway *PPARα/RXRα* activation shown at the cellular level. Grey colored molecules represent genes/proteins from our dataset.

It is expected that the number of obesity-associated loci will increase; therefore there is also a need to develop new bioinformatics tools for collecting and presenting a large amount of obesity-associated information. An option for updating our central obesity database with new loci associated with fat deposition, including miRNAs and other non-coding RNA genes, and their regulatory mechanisms will allow development of novel biomarkers and will lead to better understanding and consequently more effective treatment and control of such obesity-related disorders both in humans and in animals.

Conclusions

This study presents an integrated resource for

obesity candidate genes and miRNAs potentially involved in obese phenotypes and currently includes 1,736 loci associated to obesity in four mammalian species. The centralized online obesity database collects dispersed data in a central location and aims to be an entry point for human and animal obesity research allowing users to retrieve and submit information, which is evaluated by curators. Systems biology approach will contribute to understanding of genetic causes for obesity and also presents a novel approach to study genetic background of complex traits.

Supplementary Material

Table S1: Obesity associated miRNAs (54 human and 57 murine) with corresponding host genes. **Table S2:**

List of 175 molecules included in lipid metabolism network. <http://www.jgenomics.com/v01p0045s1.pdf>

Acknowledgements

This work was supported by the Slovenian Research Agency (ARRS) through the Research programme Comparative genomics and genome biodiversity (P4-0220).

Conflict of Interest

The authors have declared that no conflict of interest exists.

References

- [1] Herrera BM, Lindgren CM. The Genetics of Obesity. *Current Diabetes Reports*, 2010;10: 498-505.
- [2] Choquet H, Meyre D. Molecular basis of obesity: current status and future prospects. *Curr Genomics*. 2011; 12(3): 154-68.
- [3] Mutch DM, Clément K. Unraveling the Genetics of Human Obesity. *PLoS Genet*. 2006; 2(12): e188.
- [4] Kunej T, Wang Z, Michal JJ, Daniels TF, Magnuson NS, Jiang Z. Functional UQCRC1 Polymorphisms Affect Promoter Activity and Body Lipid Accumulation. *Obesity*. 2007; 15(12): 2896-901.
- [5] Jiang Z, Pappu SS, Rothschild MF. Hitting the jackpot twice: identifying and patenting gene tests related to muscle lipid accumulation for meat quality in animals and type 2 diabetes/obesity in humans. *Recent Pat DNA Gene Seq*. 2007; 1(2): 100-11.
- [6] Elías Calles JA, Gaskins CT, Busboom JR, Duckett SK, Cronrath JD, Reeves JJ, Wright RW. Differences among Wagyu sires for USDA carcass traits and palatability attributes of cooked ribeye steaks. *J Anim Sci*. 2000; 78(7): 1710-5.
- [7] Jiang Z, Kunej T, Michal JJ, Gaskins CT, Reeves JJ, Busboom JR, Dovc P, Wright RW. Significant associations of the mitochondrial transcription factor A promoter polymorphisms with marbling and subcutaneous fat depth in Wagyu x Limousin F2 crosses. *Biochem Biophys Res Commun*. 2005; 334(2): 516-23.
- [8] Rankinen T, Zuberi A, Chagnon YC, Weisnagel SJ, Argyropoulos G, Walts B, Pérusse L, Bouchard C. The human obesity gene map: the 2005 update. *Obesity (Silver Spring)*. 2006; 14(4): 529-644.
- [9] Rothschild MF, Hu ZL, Jiang Z. Advances in QTL mapping in pigs. *Int J Biol Sci*. 2007; 3(3): 192-7.
- [10] Jiang Z, Michal JJ, Williams GA, Daniels TF, Kunej T. Cross species association examination of UCN3 and CRHR2 as potential pharmacological targets for antiobesity drugs. *PLoS One*. 2006; 1:e80.
- [11] Jiang Z, Michal JJ, Chen J, Daniels TF, Kunej T, Garcia MD, Gaskins CT, Busboom JR, Alexander LJ, Wright RW, Macneil MD. Discovery of novel genetic networks associated with 19 economically important traits in beef cattle. *Int J Biol Sci*. 2009; 5(6): 528-42.
- [12] Romao JM, Jin W, Dodson MV, Hausman GJ, Moore SS, Guan IL. MicroRNA regulation in mammalian adipogenesis. *Exp Biol Med (Maywood)*. 2011; 236(9): 997-1004.
- [13] Bao B, Wang Z, Li Y, Kong D, Ali S, Banerjee S, Ahmad A, Sarkar FH. The complexities of obesity and diabetes with the development and progression of pancreatic cancer. *Biochim Biophys Acta*. 2011; 1815(2): 135-46.
- [14] Esau C, Kang X, Peralta E, Hanson E, Marcusson EG, Ravichandran LV, Sun Y, Koo S, Perera RJ, Jain R, Freier SM, Bennett CF, Lollo B, Griffey R. MicroRNA-143 regulates adipocyte differentiation. *J Biol Chem*. 2004; 279(50): 52361-5.
- [15] Georges M, Coppieters W, Charlier C. Polymorphic miRNA-mediated gene regulation: contribution to phenotypic variation and disease. *Curr Opin Genet Dev*. 2007 Jun;17(3):166-76.
- [16] Kajimoto K, Naraba H, Iwai N. MicroRNA and 3T3-L1 pre-adipocyte differentiation. *RNA*. 2006;12(9): 1626-32.
- [17] Klötting N, Berthold S, Kovacs P, Schön MR, Fasshauer M, Ruschke K, Stumvoll M, Blüher M. MicroRNA expression in human omental and subcutaneous adipose tissue. *PLoS One*. 2009; 4(3): e4699.
- [18] Mack GS. MicroRNA gets down to business. *Nat Biotechnol*. 2007; 25(6): 631-8.
- [19] Lin Q, Gao Z, Alarcon RM, Ye J, Yun Z. A role of miR-27 in the regulation of adipogenesis. *FEBS J*. 2009; 276(8): 2348-58.
- [20] Nakanishi N, Nakagawa Y, Tokushige N, Aoki N, Matsuzaka T, Ishii K, Yahagi N, Kobayashi K, Yatoh S, Takahashi A, Suzuki H, Urayama O, Yamada N, Shimano H. The up-regulation of microRNA-335 is associated with lipid metabolism in liver and white adipose tissue of genetically obese mice. *Biochem Biophys Res Commun*. 2009; 385(4): 492-6.
- [21] Ortega FJ, Moreno-Navarrete JM, Pardo G, Sabater M, Hummel M, Ferrer A, Rodriguez-Hermosa JJ, Ruiz B, Ricart W, Peral B, Fernández-Real JM. MiRNA expression profile of human subcutaneous adipose and during adipocyte differentiation. *PLoS One*. 2010; 5(2): e9022.
- [22] Takanabe R, Ono K, Abe Y, Takaya T, Horie T, Wada H, Kita T, Satoh N, Shimatsu A, Hasegawa K. Up-regulated expression of microRNA-143 in association with obesity in adipose tissue of mice fed high-fat diet. *Biochem Biophys Res Commun*. 2008; 376(4): 728-32.
- [23] Wang Q, Wang Y, Minto AW, Wang J, Shi Q, Li X, Quigg RJ. MicroRNA-377 is up-regulated and can lead to increased fibronectin production in diabetic nephropathy. *FASEB J*. 2008; 22(12): 4126-35.
- [24] Zhao E, Keller MP, Rabaglia ME, Oler AT, Stapleton DS, Schueler KL, Neto EC, Moon JY, Wang P, Wang IM, Lum PY, Ivanovska I, Cleary M, Greenawald D, Tsang J, Choi YJ, Kleinhanz R, Shang J, Zhou YP, Howard AD, Zhang BB, Kendzierski C, Thornberry NA, Yandell BS, Schadt EE, Attie AD. Obesity and genetics regulate microRNAs in islets, liver, and adipose of diabetic mice. *Mamm Genome*. 2009; 20(8): 476-85.
- [25] Kunej T, Jevsinek Skok D, Horvat S, Dovc P, Jiang Z. The glypican 3-hosted murine mir717 gene: sequence conservation, seed region polymorphisms and putative targets. *International journal of biological sciences*. 2010; 6(7): 769-72.
- [26] Jin W, Dodson MV, Moore SS, Basarab JA, Guan LL. Characterization of microRNA expression in bovine adipose tissues: a potential regulatory mechanism of subcutaneous adipose tissue development. *BMC Mol Biol*. 2010; 11: 29.
- [27] Martinelli R, Nardelli C, Pilone V, Buonomo T, Liguori R, Castano I, Buono P, Masone S, Persico G, Forestieri P, Pastore L, Sacchetti L. miR-519d Overexpression Is Associated With Human Obesity. *Obesity*. 2010 Nov;18(11):2170-6.
- [28] Twigger SN, Shimoyama M, Bromberg S, Kwitek AE, Jacob HJ, the RGD Team. The Rat Genome Database, update 2007—Easing the path from disease to data and back again. *Nucleic Acids Research*. 2007; 35(suppl 1): D658-D62.
- [29] Wang T, Li M, Guan J, Li P, Wang H, Guo Y, Shuai S, Li X. MicroRNAs miR-27a and miR-143 Regulate Porcine Adipocyte Lipid Metabolism. *International Journal of Molecular Sciences*. 2011; 12(11): 7950-9.
- [30] Baskerville S, Bartel DP. Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes. *RNA*. 2005; 11(3): 241-7.

- [31] Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004; 116(2): 281-97.
- [32] Li SC, Tang P, Lin WC. Intronic microRNA: discovery and biological implications. *DNA Cell Biol*. 2007; 26(4): 195-207.
- [33] Ogorevc J, Kunej T, Razpet A, Dovc P. Database of cattle candidate genes and genetic markers for milk production and mastitis. *Anim Genet*. 2009; 40(6): 832-51.
- [34] Ogorevc J, Dovc P, Kunej T. Comparative genomics approach to identify candidate genetic loci for male fertility. *Reprod Domest Anim*. 2011; 46(2): 229-39.
- [35] Haemmerle G, Moustafa T, Woelkart G, Büttner S, Schmidt A, van de Weijer T, Hesselink M, Jaeger D, Kienesberger PC, Zierler K, Schreiber R, Eichmann T, Kolb D, Kotzbeck P, Schweiger M, Kumari M, Eder S, Schoiswohl G, Wongsiriroj N, Pollak NM, Radner FP, Preiss-Landl K, Kolbe T, Rüllicke T, Pieske B, Trauner M, Lass A, Zimmermann R, Hoeffler G, Cinti S, Kershaw EE, Schrauwen P, Madeo F, Mayer B, Zechner R. ATGL-mediated fat catabolism regulates cardiac mitochondrial function via PPAR- α and PGC-1. *Nat Med*. 2011; 17(9): 1076-85.
- [36] Plutzky J. The PPAR-RXR Transcriptional Complex in the Vasculature. *Circulation Research*. 2011; 108(8): 1002-16.
- [37] Zorc M, Jevsinek Skok D, Godnic I, Calin GA, Horvat S, Jiang Z, Dovc P, Kunej T. Catalog of microRNA seed polymorphisms in vertebrates. *PLoS One*. 2012; 7(1): e30737.

Author Biography

Dr. Tanja Kunej is an assistant professor at the Department of Animal Science, Biotechnical Faculty, University of Ljubljana, Slovenia. Dr. Kunej is co-author of over 30 publications and has received two US patent awards. Her main research interest is integratomics in agriculture and biomedicine.

Dasa Jevsinek Skok is a PhD student of Bioscience - Bioinformatics on Biotechnical Faculty, University of Ljubljana, Slovenia. She graduated from Animal Science at the same University. She is employed as a researcher at the Department of Animal Science, Unit for pig production. Her main research interests are bioinformatics and genetics.

Minja Zorc is a PhD student of Biomedicine, University of Ljubljana, Slovenia. She graduated from Computer and information science at the same University. Her main research interest is integratomics. She is employed as a bioinformatics scientist at Rothamsted Research (BBSRC) Harpenden, UK.

Ana Ogrinc is an MSc student of Biotechnology, University of Ljubljana, Slovenia. She graduated from Biotechnology at the same University.

Jennifer J. Michal is an Associate in Research in the Department of Animal Sciences at Washington State University. She received her M.S. in Animal Sciences from the same University. She is co-author of over 40 publications.

Dr. Milena Kovac is a Professor at the Department of Animal Science, Unit for pig production, Biotechnical Faculty, University of Ljubljana, Slovenia. Dr. Kovac's main research interests are statistical methods and software development for genetic eval-

uation in livestock (PEST, PeRun, VCE), applications of genetic evaluation in livestock, pig breeding and husbandry.

Dr. Zhihua Jiang is an Associate Professor in the Animal Sciences Department at Washington State University. Since the early 1990's, Dr. Jiang's research has focused on comparative genome biology with aims at understanding how orthologous genes have evolved and shuffled during evolution. The overall objective of his research program is to develop comparative genomic tools and reagents for determining gene sequence, location, expression and function, thus advancing genome sciences and their applications in agriculture and biomedicine. Dr. Jiang has published more than 110 publications that were well cited in the field. He was editor of the book "Reproductive Genomics in Domestic Animals" Published by Wiley-Blackwell in 2010. Dr. Jiang has received eight US patent awards.